

The incidence of asymmetrical left/right skeletal and dental development in an Australian population and the effect of this on forensic age estimations

Richard B. Bassed · Christopher Briggs ·
Olaf H. Drummer

Received: 10 July 2011 / Accepted: 7 September 2011
© Springer-Verlag 2011

Abstract The prevalence of developmental asymmetry between left and right sides of the body in the third molar tooth and medial clavicular epiphysis is examined in a contemporary Australian population (92% Caucasian). The contention that differences between left and right side developmental timing is statistically insignificant, and can therefore be ignored in forensic age estimation procedures, is questioned. It was found that of a population sample of 604 individuals, 177 displayed asymmetrical timing in development between antimeres of the third molar, the medial clavicle or both. There was no correlation found between the third molar tooth and medial clavicular epiphysis in terms of left/right synchronicity. For those individuals differing in development by two or more developmental stages in either age marker or one stage in both age markers, the effect upon the accuracy of forensic age estimations can be significant. Differences in age estimates for each side were as much as 3.1 years. Age estimations based on one side only may not provide the best estimate for an individual, and more accurate results can be achieved if both sides are taken into consideration. A protocol for dealing with asymmetrical development is discussed with reference to the multifactorial age estimation method proposed by the same authors in previous research.

Keywords Forensic odontology · Forensic anthropology · Age estimation · Developmental asymmetry · CT imaging

Introduction

The development of the body's hard tissues, the skeleton and the dentition, and the changes associated with increasing chronological age, is a tool utilised in forensic practice for age determination of unknown individuals, both living and deceased. Much productive research has been conducted in this field. Accepted standards have been developed [1–6], and these standards are continually being refined and improved to make them applicable to a wider range of populations and situations [7–15].

In recent years, increased attention has been devoted to the development of the medial clavicular epiphysis and the third molar tooth, specifically to improve forensic age estimations in the hitherto problematic age cohort of the late teenage years and early adulthood (15–25 years) [9, 11, 13, 16]. This age group is difficult to assess as developing skeletal and dental sites beyond the age of 14–15 years are few. Research into skeletal and dental development of this age cohort has taken on increasing importance due to the requirement by judicial authorities to assess, in living individuals, the likelihood of having reached legal adulthood. This is especially relevant in cases of asylum seekers crossing international borders without supporting identification documentation [11, 16–19] and also in disaster victim identification where accurate age estimation can significantly aid the identification process [20].

Constructing an age estimate for an unknown individual or set of human remains involves assessing the degree of development of the age marker of interest, comparing this with published standards, preferably those having been

R. B. Bassed (✉) · C. Briggs · O. H. Drummer
Victorian Institute of Forensic Medicine and the Department of
Forensic Medicine, Monash University,
57-83 Kavanagh St, Southbank,
Melbourne, Australia
e-mail: richardb@vifm.org

C. Briggs
Department of Anatomy & Cell Biology,
University of Melbourne,
Melbourne, Australia

derived from the same or similar population as the target individual, and determining an approximate age based upon these standards. Sources of error in age estimation can be many and include utilising population standards which are not applicable to the target individual, the presence of unrecognised developmental anomalies which confound the age estimate, and finally, individual genetic variation.

A constant theme running through the literature is the oft stated conclusion that left/right developmental asymmetry is not statistically significant within a population sample and can therefore be ignored [21–31]. This leads to the conclusion that it is only necessary to evaluate one side of the body to assign an age estimate to a particular individual [3, 32–39]. Several authors have attempted to address the issue of right/left asymmetry by stating that both left and right third molars and/or clavicles need to be assessed, and either all data included into an age estimate, or either the least or most developed side used, depending upon the judicially conservative point of view relevant for that particular case [11, 13, 17, 40].

Asymmetrical development between contralateral dental and skeletal elements, also known as fluctuating asymmetry, has been recognised for many years as a normal feature of human growth, and population-based development studies have reported upon its general characteristics [41–45]. This asymmetry may result from factors such as random local genetic variation or external environmental disturbances [46]. Fluctuating asymmetry does not favour one side over the other, but can be described in terms of a normal distribution fluctuating about a mean of zero. The normally distributed nature of left/right skeletal asymmetry means that when statistical inferences are made about the development of teeth or bones on a population basis, then asymmetry becomes statistically insignificant, despite the fact that there may well be many individuals within that population who have markedly differing development timing between left and right skeletal or dental antimeres.

To date, no studies have assessed the effect of left/right developmental asymmetry upon forensic age estimations. This paper attempts to address this issue, extending previous work by the same authors on age estimation using the medial clavicular epiphysis and the third molar tooth [47–49]. The same population data used in the previous studies, based upon a post-mortem sample of the modern Australian population, were used in the current study.

Materials and methods

The individuals comprising this study were a sample of the modern Australian population, which is predominantly Caucasian (92%). The sample consisted of post-mortem high-resolution multi-detector computed tomography scans of 604 individuals in the age range of 15–25 years, gathered during medicolegal death investigation procedures carried out at the Victorian Institute of Forensic Medicine, Melbourne, Australia. The scans were conducted between January 2006 and September 2009.

The imaging system utilised was a Toshiba Aquilion 16® multi-detector scanner, which captures 1-mm thick slices of the head and neck and 2-mm slices of the remainder of the body, which are then viewed using the TeraRecon Inc Aquarius-Net® software package. This imaging protocol results in clavicular epiphyses being captured with either a 2- or 1-mm slice scan, depending upon the position of the body during the scanning process. Third molar teeth were imaged using the 1-mm protocol. Clavicles were examined using the bone algorithm, both axial and coronal slices were evaluated, and the slice displaying the greatest degree of development was scored. Third molar teeth were viewed using the Maximum Intensity Projection 3D reconstruction algorithm in the bone window, which provides what is in effect a 3D




Fig. 1 Examples of L/R developmental asymmetry in the same individual

Table 1 Numbers of cases exhibiting asymmetric left/right development

Sex	Total sample	Asymmetric molar development	Asymmetric clavicle development	Total asymmetry cases
Male	420	48 (11.4%)	84 (20%)	130 (31%)
Female	184	27 (14.6%)	24 (13%)	47 (25.5%)
Total	604	75 (12.4%)	108 (17.9%)	177 (29.3%)

The final column displaying the total number of individuals with asymmetrical development is not a sum of the molar and clavicle columns, as six individuals display asymmetry in both third molar and medial clavicle

radiograph which can be rotated in space to clearly expose the long axis of the developing tooth root.

Only those individuals who possessed both left and right antimeres for third molar teeth and medial clavicular epiphyses were included in the study. Images were examined and numerically scored based on the root lengths of the third molar teeth and the appearance and fusion status of the medial clavicular epiphysis. Third molars were scored according to the protocol developed by Demirjian [1], with the letters A–H being converted to the numbers 1–8 for ease of statistical analysis, and clavicles were scored according to Schulz et al. [29].

The sample was assessed to determine the number of cases displaying discrepant development between left and right sides, and the identified asymmetric cases were extracted and form the basis of this work. Development scores were converted into age ranges separately for left and right sides using the multiple regression formulae developed by the same authors [49]. The formulae used for males are:

$$\text{Lower age limit} = 9.91 + (0.74 \times \text{molar score}) + (1.19 \times \text{clavicle score})$$

$$\text{Upper age limit} = 11.55 + (0.99 \times \text{molar score}) + (1.41 \times \text{clavicle score})$$

and for females:

$$\text{Lower age limit} = 10.90 + (0.35 \times \text{molar score}) + (1.51 \times \text{clavicle score})$$

$$\text{Upper age limit} = 12.62 + (0.67 \times \text{molar score}) + (1.85 \times \text{clavicle score}).$$

The mean of each age range was then calculated and used as the basis for comparing differences between left

and right sides in each individual. Males and females were treated as separate groups.

Statistical analysis was performed to quantify the difference between each side and to assess how different third molar and medial clavicular development needed to be in order to have a significant effect upon calculated age estimates for individual cases. Statistical analysis was conducted using SPSS 17[®] for Windows, IBM, Chicago, IL.

Results

The sample of deceased individuals in the age range of 15–25 years consisted of 420 males and 184 females. An example of CT images depicting asymmetric development in both third molar and medial clavicle is depicted in Fig. 1.

The developmental asymmetry of the sample was assessed by independently scoring the left and right sides for both clavicle and the third molar, and then determining the degree of asymmetric development in either or both age markers for each individual. One hundred seventy-seven individuals (130 males and 47 females) displayed asymmetrical development in either or both third molar and clavicle (Table 1). The sample was further divided into those individuals differing in development by one stage only for either third molar or medial clavicle, those differing by more than one stage in either age markers, or those differing in development score in both age markers (Table 2). The mean difference in age estimate between left and right sides for the 177 asymmetric cases, males and females combined, was 1.23 years.

The difference between left and right sides in all asymmetric cases was assessed using a paired samples *t* test, males and females being assessed separately. The results of this test showed that for males, the developmental difference between left and right sides was statistically

Table 2 Breakdown of asymmetric cases by site and degree

Sex/no	1-stage difference third molar	1-stage difference clavicle	2-stage difference third molar	2-stage difference clavicle	Both sites asymmetric
Male (130)	45 (34.6%)	75 (57.7%)	4 (3.1%)	9 (6.9%)	3 (2.3%)
Female (47)	25 (53.2%)	25 (53.2%)	2 (4.2%)	0	3 (6.4%)
Total (177)	70 (39.5%)	100 (56.5%)	6 (3.4%)	9 (5.1%)	6 (3.4%)

Table 3 Typical examples of age ranges and degree of left/right discrepancy

Stage discrepancy	Sex	Age range (left)	Average (left)	Age range (right)	Average (right)	L/R discrepancy (years)
1 stage (molar)	M	15.54–18.9	17.2	14.8–17.9	16.3	0.9
	F	15.3–19.00	17.1	14.9–18.3	16.6	0.5
2 stage (molar)	M	20.6–25.1	22.8	19.1–23.1	21.1	1.7
	F	15.7–19.7	17.7	16.4–21.0	18.7	1.0
1 stage (clavicle)	M	16.3–19.9	18.1	17.5–21.3	19.4	1.3
	F	16.0–20.3	18.1	14.5–18.5	16.5	1.6
2 stage (clavicle)	M	19.4–23.7	21.5	16.0–20.9	18.4	3.1
1 stage (both)	M	18.7–22.7	20.7	16.7–20.3	18.5	2.2
	F	17.5–22.2	19.8	15.7–19.7	17.7	2.1

There were no female cases in the sample with a two-stage discrepancy between left and right medial clavicular epiphyses. Age ranges calculated using the method outlined by Bassed et al. [49]

significant ($t=2.08$, $p<0.05$), with a mean paired difference for the sample of 0.25 years. For females, the difference was not significant ($p>0.05$), and the mean paired difference for the sample was only 0.1 years.

An example of calculated age ranges (at 95% confidence interval), average ages and the differences noted between left and right sides is displayed in Table 3. It can be seen that when there is only one-stage difference in the development for either third molar or clavicle, then the resultant average discrepancy between the left and right sides is relatively minor, whereas a two-stage difference, or a difference in both molar and clavicle in the same individual, results in a much larger discrepancy. There were 21 individuals who displayed a greater than one-stage difference in third molar and/or medial clavicular development, and this accounts for 3.4% of the entire sample of 604 individuals. The differences in age estimates between the left and right sides for these individuals range from 1.0 year to a maximum of 3.1 years. The order of the effect of asymmetry in each age marker on the variation in age estimate, from greatest to least, is as follows:

1. Two-stage difference in L/R medial clavicle,
2. One-stage difference in both L/R third molar and L/R medial clavicle,

3. Two-stage difference in L/R third molar,
4. One-stage difference in L/R medial clavicle, and
5. One-stage difference in L/R third molar.

For those cases exhibiting a two-stage difference in either third molar or clavicular development, or a one-stage difference in each age marker, separate age range estimates were calculated for each side, then averaged, and the average age range compared to the actual age. An example of these cases is displayed in Table 4. Of the ten cases shown, only one has a calculated average age range that does not include the actual age of the individual concerned, being 3 months younger than the estimated lower age limit.

There was no correlation found between the third molar tooth and medial clavicular epiphysis in terms of left/right synchronicity. That is, if a left third molar was less developed than its enantiomer, there was no corresponding tendency for the medial clavicle to follow the same developmental trend.

Discussion

In forensic age estimation, practitioners attempt to derive the approximate age of an individual by applying skeletal and dental developmental data obtained from an analysis of

Table 4 L/R estimates and average age ranges compared to actual age (in years)

Gender	Age range (L)	Age range (R)	Average range	Actual age
F	17.0–21.2	15.7–19.7	16.3–20.4	16.0
M	17.5–21.3	16.0–19.3	16.7–20.3	18.2
F	15.7–19.7	16.4–21.0	16.0–20.3	18.5
F	17.5–22.2	15.7–19.7	16.6–20.9	18.7
F	16.4–21.0	18.2–23.5	17.3–22.2	18.9
M	18.7–22.7	16.7–20.3	17.7–21.5	19.7
M	16.7–20.3	18.2–22.3	17.4–21.3	20.5
M	19.4–23.7	16.0–20.9	17.7–22.3	20.9
M	16.0–20.9	17.9–21.7	16.9–21.3	21.3
F	20.9–26.6	19.7–25.4	20.3–26.0	24.3

Age ranges calculated using the method outlined by Bassed et al. [49]

a large sample of the population from which that individual comes. Inherent genetic and/or environmental variation will necessarily make any age estimation subject to error, hence the need to include an error term with each estimate. In the interest of scientific precision, this error term should be made as small as is scientifically reasonable. This is achieved by defining as accurately as possible the population parameters of the target individual and applying population data which fits most accurately with that individual. The techniques to refine population data include separating samples on the basis of gender, socioeconomic status, ancestry and geographic location [50–56]. For many unknown individuals, especially in the case of severely decomposed or skeletonised remains, this narrowing of population parameters will be practically impossible, but where it can be done with some certainty it should be attempted. It is our contention that the presence of right/left developmental asymmetry also needs to be assessed and dealt with in order to present the most accurate estimate possible with the lowest realistic error.

Fluctuating asymmetry has no bias toward either side for both medial clavicle and the third molar, and therefore any asymmetry becomes statistically invisible when a large enough sample is assessed. When all of the identified asymmetric cases are extracted and assessed separately, however, it is seen that asymmetric development is an issue on an individual basis. The significance of calculated discrepancies of up to 3.1 years between left and right sides in certain individuals implies that age estimations based on one side only may not provide the best estimate for an individual, and more accurate results can be achieved if both sides are taken into consideration.

This study has demonstrated that when there is a discrepancy of more than one developmental stage between antimeres in either the third molar or medial clavicular epiphysis, then the difference in age estimate for left and right sides can be significant. Discrepancies in clavicular development have a greater effect on the resultant age estimate than third molar teeth, with the greatest effect seen in those individuals differing by two developmental stages for left and right clavicular epiphyses. Although the percentage of individuals with greater than one-stage difference between left and right sides is small (3.4%), the fact remains that in order for age estimates to be as precise and as accurate as possible for the greatest number of individuals, then it is necessary to develop a protocol to deal with this fluctuating asymmetry.

It is proposed that, where possible, both left and right sides be assessed for any anatomical age marker being examined. If it is found that there is one stage or less discrepancy, then the age estimate can be calculated using whichever side is the most judicially conservative. In these cases, the discrepancy in the age range for each side will

not be significant enough to result in any meaningful inaccuracy. If, however, there is greater than one developmental stage discrepancy between antimeres, then it is recommended that age ranges be calculated individually for each side and the average of these calculations used as the final age range. An example of this is depicted in Table 4, which shows that in nine out of the ten cases presented, the actual age of the individual was within the calculated average age range, with the remaining case being 3 months younger than the age range would suggest. If the single side age ranges were to be used, then in three of the cases the actual age would have fallen outside the calculated range, in one case by as much as 1 year.

In summary, this research has shown that whilst fluctuating asymmetry occurring in development timing of the medial clavicular epiphysis and the third molar tooth may be insignificant on a population-wide basis, when assessing individual forensic cases, these developmental left/right discrepancies need to be considered where the difference in developmental staging is greater than one stage. A method of averaging the age ranges for each side is presented which should improve the accuracy of forensic age estimation procedures.

References

- Demirjian A, Goldstein H, Tanner JM (1973) A new system of dental age assessment. *Hum Biol* 45(2):211–227
- Mckern TW, Stewart TD (1957) Skeletal age changes in young American males analysed from the standpoint of identification. Natick, Massachusetts, Headquarters, Quartermaster Research & Development Command
- Webb PA, Suchey JM (1985) Epiphyseal union of the anterior iliac crest and medial clavicle in a modern multiracial sample of American males and females. *Am J Phys Anthropol* 68(4):457–466
- Greulich WW, Pyle SI (1959) Radiographic atlas of skeletal development of the hand and wrist, 2nd edn. Stanford University Press, California
- Tanner JM, Whitehouse RH, Marshall WA, Healy MJR, Goldstein H (1975) Assessment of skeletal maturity and prediction of adult height (tw2 method). Academic, New York
- Moorrees CF, Fanning EA, Hunt EE Jr (1963) Age variation of formation stages for ten permanent teeth. *J Dent Res* 42:1490–1502
- Olze A, Taniguchi M, Schmeling A, Zhu BL, Yamada Y, Maeda H et al (2003) Comparative study on the chronology of third molar mineralization in a Japanese and a German population. *Leg Med (Tokyo)* 5(Suppl 1):S256–S260
- Orhan K, Ozer L, Orhan AI, Dogan S, Paksoy CS (2007) Radiographic evaluation of third molar development in relation to chronological age among Turkish children and youth. *Forensic Sci Int* 165(1):46–51
- Cardoso HF (2008) Age estimation of adolescent and young adult male and female skeletons II, epiphyseal union at the upper limb and scapular girdle in a modern Portuguese skeletal sample. *Am J Phys Anthropol* 137(1):97–105
- Cardoso HF (2007) Environmental effects on skeletal versus dental development: using a documented subadult skeletal sample

- to test a basic assumption in human osteological research. *Am J Phys Anthropol* 132(2):223–233
11. Kellinghaus M, Schulz R, Vieth V, Schmidt S, Schmeling A (2010) Forensic age estimation in living subjects based on the ossification status of the medial clavicular epiphysis as revealed by thin-slice multidetector computed tomography. *Int J Legal Med* 124(2):149–154
 12. Schulz R, Muhler M, Reisinger W, Schmidt S, Schmeling A (2008) Radiographic staging of ossification of the medial clavicular epiphysis. *Int J Legal Med* 122(1):55–58
 13. Kasper KA, Austin D, Kvanli AH, Rios TR, Senn DR (2009) Reliability of third molar development for age estimation in a Texas Hispanic population: a comparison study. *J Forensic Sci* 54(3):651–657
 14. Blenkin MR, Evans W (2011) Age estimation from the teeth using a modified Demirjian system. *J Forensic Sci* 55(6):1504–1508
 15. Graham JP, O'Donnell CJ, Craig PJ, Walker GL, Hill AJ, Cirillo GN et al (2010) The application of computerized tomography (CT) to the dental ageing of children and adolescents. *Forensic Sci Int* 195(1–3):58–62
 16. Schmeling A, Grundmann C, Fuhrmann A, Kaatsch HJ, Knell B, Ramsthaler F et al (2008) Criteria for age estimation in living individuals. *Int J Legal Med* 122(6):457–460
 17. Mincer HH, Harris EF, Berryman HE (1993) The A.B.F.O. study of third molar development and its use as an estimator of chronological age. *J Forensic Sci* 38(2):379–390
 18. Cunha E, Baccino E, Martrille L, Ramsthaler F, Prieto J, Schuliar Y et al (2009) The problem of aging human remains and living individuals: a review. *Forensic Sci Int* 193(1–3):1–13
 19. Garamendi PM, Landa MI, Ballesteros J, Solano MA (2005) Reliability of the methods applied to assess age minority in living subjects around 18 years old. A survey on a Moroccan origin population. *Forensic Sci Int* 154(1):3–12
 20. Bassed RB, Hill AJ (2011) The use of computed tomography (CT) to estimate age in the 2009 Victorian bushfire victims: a case report. *Forensic Sci Int* 205(1–3):48–51
 21. Johanson G (1971) Age determinations from human teeth, a critical evaluation with special consideration of changes after fourteen years of age (Odontol revy v.22. supplement). Glerup, Allingabro
 22. Bhat VJ, Kamath GP (2007) Age estimation from root development of mandibular third molars in comparison with skeletal age of wrist joint. *Am J Forensic Med Pathol* 28(3):238–241
 23. Meisl A, Tangl S, Huber C, Maurer B, Watzek G (2007) The chronology of third molar mineralization in the Austrian population—a contribution to forensic age estimation. *Forensic Sci Int* 169(2–3):161–167
 24. Prieto JL, Barberia E, Ortega R, Magana C (2005) Evaluation of chronological age based on third molar development in the Spanish population. *Int J Legal Med* 119(6):349–354
 25. Kullman L, Martinsson T, Zimmerman M, Welander U (1995) Computerized measurements of the lower third molar related to chronologic age in young adults. *Acta Odontol Scand* 53(4):211–216
 26. Kullman L, Johanson G, Akesson L (1992) Root development of the lower third molar and its relation to chronological age. *Swed Dent J* 16(4):161–167
 27. Willershausen B, Löffler N, Schulze R (2001) Analysis of 1202 orthopantomograms to evaluate the potential of forensic age determination based on third molar developmental stages. *Eur J Med Res* 6(9):377–384
 28. Thevissen PW, Pittayapat P, Fieuws S, Willems G (2009) Estimating age of majority on third molars developmental stages in young adults from Thailand using a modified scoring technique. *J Forensic Sci* 54(2):428–432
 29. Schulz R, Muhler M, Mutze S, Schmidt S, Reisinger W, Schmeling A (2005) Studies on the time frame for ossification of the medial epiphysis of the clavicle as revealed by ct scans. *Int J Legal Med* 119(3):142–145
 30. Kreitner KF, Schweden FJ, Riepert T, Nafe B, Thelen M (1998) Bone age determination based on the study of the medial extremity of the clavicle. *Eur Radiol* 8(7):1116–1122
 31. Schmeling A, Schulz R, Reisinger W, Muhler M, Wernecke KD, Geserick G (2004) Studies on the time frame for ossification of the medial clavicular epiphyseal cartilage in conventional radiography. *Int J Legal Med* 118(1):5–8
 32. De Salvia A, Calzetta C, Orrico M, De Leo D (2004) Third mandibular molar radiological development as an indicator of chronological age in a European population. *Forensic Sci Int* 146(Suppl):S9–S12
 33. Gorgani N, Sullivan RE, DuBois L (1990) A radiographic investigation of third-molar development. *ASDC J Dent Child* 57(2):106–110
 34. Levesque GY, Demirjian A, Tanguay R (1981) Sexual dimorphism in the development, emergence, and agenesis of the mandibular third molar. *J Dent Res* 60(10):1735–1741
 35. Nortje CJ (1983) The permanent mandibular third molar. Its value in age determination. *J Forensic Odontostomatol* 1(1):27–31
 36. Olze A, Bilang D, Schmidt S, Wernecke KD, Geserick G, Schmeling A (2005) Validation of common classification systems for assessing the mineralization of third molars. *Int J Legal Med* 119(1):22–26
 37. Schulz R, Zwiesigk P, Schiborr M, Schmidt S, Schmeling A (2008) Ultrasound studies on the time course of clavicular ossification. *Int J Legal Med* 122(2):163–167
 38. Mesotten K, Gunst K, Carbonez A, Willems G (2003) Chronological age determination based on the root development of a single third molar: a retrospective study based on 2513 OPGs. *J Forensic Odontostomatol* 21(2):31–35
 39. Cameriere R, Ferrante L, De Angelis D, Scarpino F, Galli F (2008) The comparison between measurement of open apices of third molars and Demirjian stages to test chronological age of over 18 year olds in living subjects. *Int J Legal Med* 122(6):493–497
 40. Knell B, Ruhstaller P, Prieels F, Schmeling A (2009) Dental age diagnostics by means of radiographical evaluation of the growth stages of lower wisdom teeth. *Int J Legal Med* 123(6):465–469
 41. Baer MJ, Dyrkatz J (1957) Bilateral asymmetry in skeletal maturation of the hand and wrist: a roentgenographic analysis. *Am J Phys Anthropol* 15(2):181–196
 42. Fazekas IG, Kosa F (1978) Forensic fetal osteology. Akademiai Kiado, Budapest
 43. Albert AM, Greene DL (1999) Bilateral asymmetry in skeletal growth and maturation as an indicator of environmental stress. *Am J Phys Anthropol* 110(3):341–349
 44. Dreizen S, Spirakis CN, Stone RE (1967) A comparison of skeletal growth and maturation in undernourished and well-nourished girls before and after menarche. *J Pediatr* 70(2):256–263
 45. Rios L, Cardoso HF (2009) Age estimation from stages of union of the vertebral epiphyses of the ribs. *Am J Phys Anthropol* 140(2):265–274
 46. Steele J (2000) Skeletal indicators of handedness. In: Cox M, Mays S (eds) *Human osteology in archaeology and forensic science*. Cambridge University Press, Cambridge, pp 307–323
 47. Bassed RB, Briggs C, Drummer OH (2011) Age estimation and the developing third molar tooth: an analysis of an Australian population using computed tomography. *J Forensic Sci* 56(5):1185–1191
 48. Bassed RB, Drummer OH, Briggs C, Valenzuela A (2011) Age estimation and the medial clavicular epiphysis: analysis of the age of majority in an Australian population using computed tomography. *Forensic Sci Med Pathol* 7(2):148–154

49. Bassed RB, Briggs C, Drummer OH (2011) Age estimation using CT imaging of the third molar tooth, the medial clavicular epiphysis, and the sphenoid-occipital synchondrosis: a multifactorial approach. *Forensic Sci Int*. doi:[10.1016/j.forsciint.2011.06.007](https://doi.org/10.1016/j.forsciint.2011.06.007)
50. Martin-de las Heras S, Garcia-Forteza P, Ortega A, Zodicovich S, Valenzuela A (2008) Third molar development according to chronological age in populations from Spanish and Magrebian origin. *Forensic Sci Int* 174(1):47–53
51. Olze A, Schmeling A, Taniguchi M, Maeda H, van Niekerk P, Wernecke KD et al (2004) Forensic age estimation in living subjects: the ethnic factor in wisdom tooth mineralization. *Int J Legal Med* 118(3):170–173
52. Olze A, van Niekerk P, Ishikawa T, Zhu BL, Schulz R, Maeda H et al (2007) Comparative study on the effect of ethnicity on wisdom tooth eruption. *Int J Legal Med* 121(6):445–448
53. Schmeling A, Olze A, Reisinger W, Geserick G (2005) Forensic age estimation and ethnicity. *Leg Med (Tokyo)* 7(2):134–137
54. Schmeling A, Olze A, Reisinger W, König M, Geserick G (2003) Statistical analysis and verification of forensic age estimation of living persons in the Institute of Legal Medicine of the Berlin University Hospital Charité. *Leg Med (Tokyo)* 5(Suppl 1):S367–S371
55. Thevissen PW, Fieuws S, Willems G (2010) Human third molars development: comparison of 9 country specific populations. *Forensic Sci Int* 201(1–3):102–105
56. Thevissen PW, Alqerban A, Asaumi J, Kahveci F, Kaur J, Kim YK et al (2010) Human dental age estimation using third molar developmental stages: accuracy of age predictions not using country specific information. *Forensic Sci Int* 201(1–3):106–111